

Caution: For Laboratory Use. A research reagent for research purposes only

Human Vasopressin V2 Receptor, γ -Irradiated Frozen Cells

Product No.: ES-363-CF

Lot No.: 1003

Material Provided

Cells:	1 x 1 mL frozen aliquot
Format:	~2.5 x 10 ⁶ cells / mL in DMEM, 10% FBS, 1 mM sodium pyruvate with 10 % DMSO

Product Information

Cellular Background:	CHO-K1
Frozen cells info:	Frozen recombinant, γ -irradiated CHO-K1 cells expressing the human Vasopressin V2 receptor.
DNA Sequence:	Identical to coding sequence of NM_000054.4 with the exception of a silent mutation in codon # 309 (CTA becomes CTG, both coding for a Leu).
Corresponding Protein Sequence:	Identical to GenBank NP_000045.1.
Storage Conditions:	Store in liquid nitrogen (vapor phase) immediately upon receipt, or maximum 15 days at -80°C. cAMPZen [®] is designed for single use only. Do not refreeze.

Quality Control

EC₅₀ for a reference agonist is determined using a TR-FRET cAMP assay (Figure 1). Mycoplasma test is performed using MycoAlert[®] Mycoplasma detection kit. We certify that these results meet our quality release criteria.


(ARG⁸)-VASOPRESSIN - (EC₅₀):	1.6 pM
Mycoplasma:	This cell line tested negative for Mycoplasma.

Recommended Use and Thawing Conditions

Assay Medium: HBSS 1X (without phenol red); 5 mM HEPES; BSA-stabilizer (7.5% solution); 0.5 mM IBMX (pH 7.4).

Thawing Cells: Using appropriate personal protective equipment, place the frozen aliquot in a 37°C water bath and agitate until its content is thawed completely. Immediately remove from water bath, transfer content to 10 mL assay medium and centrifuge (150 x g, 5 min). Resuspend cell pellet in appropriate volume of assay medium for use in the selected LANCE[®] cAMP assay (see LANCE[®] cAMP Assays section).


Typical Product Data



Agonist	EC ₅₀ (M)
(Arg ⁸)-Vasopressin	8.9 x 10 ⁻¹²
(deamino-Cys ¹ , Val ⁴ , D-Arg ⁸)-Vasopressin	2.0 x 10 ⁻¹²
Oxytocin	1.0 x 10 ⁻⁹

Figure 1: Agonist Response in TR-FRET cAMP assay

An agonist dose-response experiment was performed in 384-well format using 1 000 cells/well. Reader: RUBYstar. Data from a representative experiment are shown. The Z'-factor was calculated for (Arg⁸)-Vasopressin with at least 16 background and 16 maximal signal points (Z' = 0.80).



Antagonist	IC ₅₀ (M)
[Adamantaneacetyl ¹ , O-Et-D-Tyr ² , Val ⁴ , Abu ⁶ , Arg ^{8,9}]-Vasopressin	1.3 x 10 ⁻⁷

Figure 2: Antagonist Response in TR-FRET cAMP assay

An antagonist dose-response experiment was performed in 384-well format using 1 000 cells/well and 0.03 nM (Arg⁸)-Vasopressin (reference agonist). Reader: RUBYstar. Data from a representative experiment are shown.

Table 1. References of compounds used for functional characterization

Name	Provider	Cat n°	Working Stock Solution
(Arg ⁸)-Vasopressin	Bachem	H-1780	1 mM in water
(deamino-Cys ¹ , Val ⁴ , D-Arg ⁸)-Vasopressin	Sigma	V-2013	1 mM in water
Oxytocin	Bachem	H-2510	10 mM in water
[Adamantaneacetyl ¹ , O-Et-D-Tyr ² , Val ⁴ , Abu ⁶ , Arg ^{8,9}]-Vasopressin	Sigma	V-2381	10 mM in water

LANCE[®] cAMP Assays

LANCE[®] Ultra cAMP - please refer to the user manual included with the LANCE[®] Ultra cAMP kits:

TRF0262 – 1 000 Assay points

TRF0263 – 10 000 Assay points

TRF0264 – 50 000 Assay points

LANCE[®] cAMP - please refer to the user manual included with the LANCE[®] cAMP kits:

AD0262 – 500 Assay points

AD0262E – 1 000 Assay points

AD0263 – 10 000 Assay points

AD0264 – 50 000 Assay points

Suggested Materials and Instrumentation

Please visit our website

www.perkinelmer.com/CellLines

This product is not for resale or distribution except by authorized distributors.

LIMITED WARRANTY: PerkinElmer BioSignal Inc. warrants that, at the time of shipment, the products sold by it are free from defects in material and workmanship and conform to specifications which accompany the product. PerkinElmer BioSignal Inc. makes no other warranty, express or implied with respect to the products, including any warranty of merchantability or fitness for any particular purpose. Notification of any breach of warranty must be made within 60 days of receipt unless otherwise provided in writing by PerkinElmer BioSignal Inc. No claim shall be honored if the customer fails to notify PerkinElmer BioSignal Inc. within the period specified. The sole and exclusive remedy of the customer for any liability of PerkinElmer BioSignal Inc. of any kind including liability based upon warranty (express or implied whether contained herein or elsewhere), strict liability contract or otherwise is limited to the replacement of the goods or the refunds of the invoice price of goods. PerkinElmer BioSignal Inc. shall not in any case be liable for special, incidental or consequential damages of any kind.

PerkinElmer, Inc.
940 Winter Street
Waltham, MA 02451 USA
P: (800) 762-4000 or
(+1) 203-925-4602
www.perkinelmer.com



For a complete listing of our global offices, visit www.perkinelmer.com/ContactUs

Copyright ©2009, PerkinElmer, Inc. All rights reserved. PerkinElmer[®] is a registered trademark of PerkinElmer, Inc. All other trademarks are the property of their respective owners.