

Research use only. Not for use in diagnostic procedures.

human Opioid delta Receptor Cell Line

Product No.: RBHODM-K

Lot No.: 2533360

Material Provided

Cells: 2 x 1 mL frozen aliquot (RBHODM-WV)
Format: ~5 x 10⁶ cells/mL in freezing medium

Product Information

Cellular Background: CHO-K1

GenBank Accession Number: U10504

Cell Line Development: Our expression plasmid containing the sequence coding for the human Opioid delta receptor was transfected in CHO-K1 cells. Hygromycin-resistant clones were obtained by limit dilution and compared for receptor expression levels by radioligand binding assay. The clone with the highest receptor expression level was selected for characterization in binding and functional assays.

Receptor expression level (B_{MAX}): Estimated to be 23 pmol/mg protein, using [³H]-Naltrindole*

Shipping Conditions: Shipped on dry ice. Please ensure dry ice is still present in the package upon receipt or contact customer support.

Storage Conditions: Store in liquid nitrogen (vapor phase) immediately upon receipt.

Quality Control

Mycoplasma: This cell line tested **negative** for mycoplasma.

Recommended Cell Culture Conditions (CHO-K1)

- The recommended media catalogue number and supplier reference information are listed in this Product Technical Data Sheet (last page). Media composition is specifically defined for each cell type and receptor expression selection. The use of incorrect media or component substitutions can lead to reduced cell viability, growth issues and/or altered receptor expression.
- Cells undergo major stress upon thawing, and need to adapt to their new environment which may initially affect cell adherence and growth rates. The initial recovery of the cells, and initial doubling time, will vary from laboratory to laboratory, reflecting differences in the origin of culture media and serum, and differences in methodology used within each laboratory.
- For the initial period of cell growth (i.e. until cells have reached Log-phase, typically 4-10 days), we strongly recommend removal of the antibiotics (G418, Zeocin™, Puromycin, Blasticidin, Hygromycin, Penicillin and Streptomycin) from the culture media. Immediately after thawing, cells may be more permeable to antibiotics, and a higher intracellular antibiotic concentration may result as a consequence. Antibiotics should be re-introduced when cells have recovered from the thawing stress.

Growth Medium: Ham's F-12, 10% Fetal Clone II, 1 mM Sodium Pyruvate, 250 µg/ml Hygromycin B

Freezing Medium: Ham's F-12, 10% Fetal Clone II with 10% DMSO, without selection agents.

Thawing Cells: Using appropriate personal protective equipment, rapidly place the frozen aliquot in a 37°C water bath (do not submerge) and agitate until its content is thawed completely. Immediately remove from water bath, spray aliquot with 70% ethanol and wipe excess. Under aseptic conditions using a sterile pipette, transfer content to a sterile centrifuge tube containing 10 mL growth medium without antibiotics, pre-warmed at 37°C, and centrifuge (150 x g, 5 min). Discard supernatant using a sterile pipette. Resuspend cell pellet in 10 mL of pre-warmed growth medium without antibiotics by pipetting up and down to break up any clumps, and transfer to an appropriate culture flask (e.g. T-25, T-75 or T-175, see recommended seeding density below). Cells are cultured as a monolayer at 37°C in a humidified atmosphere with 5% CO₂.

Recommended Seeding Density: Thawing: 15 000 – 33 000 cells/cm²
Log-phase: 11 000 – 15 000 cells/cm²

Troubleshooting: Initial doubling time can vary between 18 and 96 hours (Average = 25 hours). If cells are still not adhering after 48 hours or grow very slowly, we recommend maintaining the cells in culture and not replacing the media before 5-6 days (cells secrete factors that can help with adherence and growth). If confluence is still <50% after 5-6 days, it is recommended that you replace the media with fresh media (without antibiotics). Do not passage the cells until they reach 80-90% confluence (Log-phase). If cells have not recovered after 10-12 days, please contact our Technical Support.

Culture Protocol: Under aseptic conditions, cells are grown to 80% confluence (Log-phase) and trypsinized (0.05% trypsin / 0.5 mM EDTA in calcium and magnesium-free PBS). See recommended seeding density for Log-phase above.

Banking Protocol: Cells are grown to 70-80% confluence (Log-phase). Under aseptic conditions, remove medium and rinse the flask with an appropriate volume of calcium and magnesium-free PBS (example 10 mL for T-175). Trypsinize (0.05% trypsin / 0.5 mM EDTA in calcium and magnesium-free PBS) to detach cells (example 5 mL for T-175), let stand 5-10 min at 37°C. Add fresh, room temperature growth medium (without antibiotics) to stop trypsinization and dilute EDTA (example 10 mL for T-175). Transfer cells to a sterile centrifuge tube and centrifuge (150 x g, 5 min). Discard supernatant using a sterile pipette. Resuspend cell pellet in ice-cold freezing medium by pipetting up and down to break up any clumps. Count cells and rapidly aliquot at the selected cell density (e.g. 2.5 x 10⁶ cells/mL) in sterile polypropylene cryovials. Use appropriate material to ensure slow cooling (about 1°C/min) until -70°C. Transfer vials into a liquid nitrogen tank (vapour phase) for storage.

Reference of Cell Culture Media

Name	Provider	Cat. Number
Ham's Nutrient Mixture F-12	Fisher Scientific	SH30026.02
DMEM/High Glucose	Fisher Scientific	SH30022.02
Advanced DMEM/F12	ThermoFisher (Gibco)	12634-010
EMEM	Lonza	06-174G
EX-CELL® CHO DHFR-	Sigma	C8862
Fetal Clone II	Fisher Hyclone	SH30066.03
FBS	RMBIO	FBS-BBT
FBS dialyzed	Wisent	080950
G418 Sulfate	Wisent	400-130-IG
Zeocin™	ThermoFisher (Gibco)	R25005
Blasticidin	ThermoFisher (Gibco)	R210-01
Puromycin	Wisent	400-160-EM
Hygromycin D	ThermoFisher Scientific	10687010
Trypsin-EDTA	Fisher Scientific	SH30236.02
Sodium Pyruvate	ThermoFisher (Gibco)	11360
L-Glutamine	ThermoFisher (Gibco)	25030

Historical Cell Line Validation using Membrane Preparation*

Saturation and Competition Binding Assay

Assay Buffer: 50 mM Tris-HCl pH 7.4, 100 mM NaCl

Wash Buffer: 50 mM Tris-HCl pH 7.4, 100 mM NaCl

Binding Protocol: Binding assays are performed in 550 μ L total volume according to the following conditions:

1 - Membrane dilution: 0.125 mL of membranes + 24.875 mL assay buffer (1:200 dilution)

2 - Incubation: 25 μ L of incubation buffer or Naltrexone (Sigma N3136) 100 μ M final for non specific binding (Saturation binding assay)

For competition binding assay: 25 μ L of reference compounds at decreasing concentrations (see figure 2)

25 μ L of radioligand at the appropriate concentration (see graph below)
500 μ L of diluted membranes

3 - Incubation time: 60 minutes at RT

4 - Filtration: aspirate and wash 9 x 500 μ L with ice cold wash buffer over GF/C filter (presoaked in 0.5 % PEI).




Figure 1: Saturation binding assay curve (filtration)

96-well saturation binding assay curve (12 μ g membranes/well, TopCount®) using [³H]-Naltrindole (PerkinElmer NET1065 Lot No.: CC80750)

Typical Product Data




Figure 2: Competition binding assay curve (filtration)

96-well competition binding assay curve (15 μ g membranes/well, TopCount®). Recommended radioligand concentration = 0.4 nM.

*Even though two sites can be observed occasionally with some ligands, the data presented is derived from single site fitting.

Reference Compounds	K _i (nM)
Naltriben	1.6
BNTX	37
U-69593	47145

* Membrane preparation available as RBHODM400UA

Please visit our website: <http://www.perkinelmer.com/category/receptor-cell-lines-membranes> for additional information on cell lines, associated reagents, membranes and instrumentation.

This product is not for resale or distribution except by authorized distributor.