

Caution: For Laboratory Use. A product for research purposes only

## **Eu-W1024-labeled Anti-phospho-Histone H3 (Thr3) Antibody**

**Product No.:** TRF0211-D / TRF0211-M

**Lot No.:** 2707916

### **Material Provided**

**Format:** TRF0211-D 10 µg (1 562 assay points\*)  
TRF0211-M 100 µg (15 625 assay points\*)  
\*Assuming 40 fmol/ assay point

**Volume:** 100 µL (TRF0211-D) or 1 mL (TFR0211-M)

**Manufacturing Date:** February 7, 2020

### **Product Information**

**Antibody:** Europium-labeled rabbit monoclonal antibody recognizing phospho-Thr3 in human Histone H3.

**Molecular Weight:** 160 000

**Storage Buffer:** 50 mM Tris-HCl (pH 7.4), 0.9% NaCl, 0.1% BSA and 0.05% sodium azide as preservative

**Stability:** This product is stable for at least **12 months** from the manufacturing date when stored in its original packaging and the recommended storage conditions.

**Storage Conditions:** Store at -20°C. Repeated freezing and thawing should be avoided.

**Safety Note:** The storage buffer contains sodium azide (NaN<sub>3</sub>) as a preservative. Disposal of all waste should be in accordance with local regulations.

### **Quality Control**

The QC release specifications are based on spectrophotometric analysis of the labeled antibody. We certify that these results meet our quality release criteria.

**Labeling Ratio:** 7.3 (Eu/Ab)

**Concentration:** 100 µg/mL (0.625 µM)

## Recommended Assay Conditions

### RSK2 kinase: ATP titration

#### Reagent Preparation:

- Prepare 1X Kinase Assay Buffer: 50 mM HEPES pH 7.5, 1 mM EGTA, 10 mM MgCl<sub>2</sub>, 2 mM DTT and 0.01% Tween-20.
- Prepare a 2X RSK2 solution: dilute enzyme to a concentration of 8 nM in 1X Kinase Assay Buffer. Keep on ice.
- Prepare a 4X *ULight*-Histone H3 (Thr3/Ser10) Peptide solution: dilute *ULight*-Histone H3 (Thr3/Ser10) Peptide to a concentration of 200 nM in 1X Kinase Assay Buffer.
- Prepare a 4X ATP solution: dilute ATP to concentrations ranging from 40 nM to 4 mM (serial half-log dilutions) in 1X Kinase Assay Buffer. Keep on ice.
- Prepare 1X Detection Buffer: dilute 0.5 mL of 10X Detection Buffer with 4.5 mL of H<sub>2</sub>O.
- Prepare a 4X Stop solution\*: dilute EDTA to a concentration of 40 mM in 1X Detection Buffer.
- Prepare a 4X Detection Mix: dilute Europium-anti-phospho-Histone H3 (Thr3) Antibody to a concentration of 8 nM in 1X Detection Buffer.


#### Protocol:

- Pipet 5 µL of 2X RSK2 solution into a 384-well white OptiPlate-384 (4 nM final concentration).
- Add 2.5 µL of 4X *ULight*-Histone H3 (Thr3/Ser10) Peptide solution (50 nM final concentration).
- Add 2.5 µL of 4X ATP solution (10 nM to 1 mM final concentrations).
- Cover plate with TopSeal-A and incubate 45 min at 23°C.
- Add 5 µL of 4X Stop solution\* and incubate 5 min at 23°C.
- Add 5 µL of 4X Detection Mix (2 nM Europium-anti-phospho-Histone H3 (Thr3) Antibody final concentration).
- Cover plate with TopSeal-A and incubate 60 min at 23°C.
- Remove TopSeal-A and read in TR-FRET mode at 665 nm (excitation at 320 or 340 nm).

*\*Alternatively, the Stop solution and Detection Mix can be premixed and added together to the kinase reaction.*

## Typical ATP Titration Data

RSK2 kinase assay using *ULight*-Histone H3 (Thr3/Ser10) Peptide and Eu-anti-phospho-Histone H3 (Thr3) Antibody obtained using the EnVision® Multilabel Reader:



## Suggested Materials

	Supplier	Cat. No.
• Substrate: <i>ULight</i> <sup>™</sup> - Histone H3 (Thr3/Ser10) Peptide	PerkinElmer	TRF0125
• Antibody: Eu- anti-phospho-Histone H3 (Thr3) Antibody	PerkinElmer	TRF0211
• Kinase: RSK2	Carna Biosciences	01-150
• Detection Buffer: LANCE <sup>®</sup> Detection Buffer, 10X	PerkinElmer	CR97-100
• Plate: OptiPlate <sup>™</sup> -384, white	PerkinElmer	6007299
• TopSeal <sup>™</sup> : TopSeal-A	PerkinElmer	6050195

Please visit our website for additional resource:

[www.perkinelmer.com/LANCE](http://www.perkinelmer.com/LANCE)

**This product is not for resale or distribution except by authorized distributors.**

LANCE<sup>®</sup> products may be covered by one or more the following patents: US 5,202,423 and US 5,324,825

**LIMITED WARRANTY:** PerkinElmer BioSignal Inc. warrants that, at the time of shipment, the products sold by it are free from defects in material and workmanship and conform to specifications which accompany the product. PerkinElmer BioSignal Inc. makes no other warranty, express or implied with respect to the products, including any warranty of merchantability or fitness for any particular purpose. Notification of any breach of warranty must be made within 60 days of receipt unless otherwise provided in writing by PerkinElmer BioSignal Inc. No claim shall be honored if the customer fails to notify PerkinElmer BioSignal Inc. within the period specified. The sole and exclusive remedy of the customer for any liability of PerkinElmer BioSignal Inc. of any kind including liability based upon warranty (express or implied whether contained herein or elsewhere), strict liability contract or otherwise is limited to the replacement of the goods or the refunds of the invoice price of goods. PerkinElmer BioSignal Inc. shall not in any case be liable for special, incidental or consequential damages of any kind.

PerkinElmer, Inc.  
940 Winter Street  
Waltham, MA 02451 USA  
P: (800) 762-4000 or  
(+1) 203-925-4602  
[www.perkinelmer.com](http://www.perkinelmer.com)



For a complete listing of our global offices, visit [www.perkinelmer.com/ContactUs](http://www.perkinelmer.com/ContactUs)

Copyright ©2009, PerkinElmer, Inc. All rights reserved. PerkinElmer<sup>®</sup> is a registered trademark of PerkinElmer, Inc. All other trademarks are the property of their respective owners.